

Finite Fields and Their Applications **8**, 511–518 (2002)

doi:10.1006/ffta.2002.0360

Reed–Muller-Type Codes Over the Segre Variety

M. González-Sarabia and C. Rentería¹*Escuela Superior de Física y Matemáticas, IPN, 07300 México, D.F., México*

E-mail: msarabia@campus.cem.itesm.mx; renteri@esfm.ipn.mx

and

H. Tapia-Recillas²*Departamento de Matemáticas, Universidad Autónoma Metropolitana-I, 09340,
México, D.F., México*

E-mail: htr@xanum.uam.mx

Communicated by James W.P. Hirschfeld

Received February 9, 2001; revised February 24, 2002; published online June 14, 2002

The a -invariant is determined and a description of the defining ideal for the set S_K of rational points of the Segre variety over a finite field K is given. The dimension as well as the minimum distance of a Reed–Muller-type linear code defined over S_K are also determined. An example is given to illustrate the ideas. © 2002 Elsevier Science (USA)

Key Words: Reed–Muller codes; Segre variety; a -invariant; defining ideal.

1. INTRODUCTION

Let $K = GF(q)$ be a finite field with $q = p^r$ elements (p prime and r a positive integer). Let $\mathbf{P}_l(K)$ be the l -projective space over K and let $X = \{P_1, \dots, P_t\}$ be a subset of $\mathbf{P}_l(K)$. If \mathcal{L} is a finite K -dimensional linear space of functions defined on the set X with values in the field K , the evaluation map

$$ev_X: \mathcal{L} \rightarrow K^t, \quad ev_X(f) = (f(P_1), \dots, f(P_t))$$

¹Partially supported by COFAA-IPN and SNI-SEP, México.

²Partially supported by CONACyT and SNI-SEP, México.



determines the K -linear code $C_X = ev_X(\mathcal{L})$. For any subset X and a general linear subspace of functions \mathcal{L} , it is a difficult task to determine the parameters of the code C_X .

Let $A = K[X_0, \dots, X_l] = \bigoplus_{i \geq 0} A_i$ be the graded polynomial ring over the field K where A_i is the K -subspace generated by all monomials in A of degree i . If $X \subseteq \mathbf{P}_l(K)$ and $\mathcal{L} = A_d$, ($d \geq 1$) is the d -graded homogeneous component of the ring A , the corresponding linear code $C_X(d) := ev_X(A_d)$, will be called the Reed–Muller linear code over the set X . If $I_X = \{f \in A : f(P) = 0 \ \forall P \in X\} = \bigoplus_{i \geq 0} I_X(i)$ is the (graded) vanishing ideal of the set X , the linear code $C_X(d)$ is isomorphic to the K -vector space $A_d/I_X(d)$. Thus, the dimension of the code is given by the *Hilbert function* of the ring A/I_X . A generating matrix for this code can be obtained by finding a Gröbner basis for the ideal I_X such that the cosets, with respect to this ideal, of the monomials in A of degree d which are not in the leading terms ideal $LT(I_X)$, are a K -basis for the vector space $A_d/I_X(d)$. If \mathcal{B} is this set of monomials then $(ev_X(\mathcal{M}))_{\mathcal{M} \in \mathcal{B}}$ is a generating matrix for the code $C_X(d)$.

Several authors have provided results about the code $C_X(d)$ or related codes for different sets X [2, 4, 8–12]. The Segre embedding has been used recently to prove a conjecture on the weight of product codes [13] proposed by Wei and Yang [15]. The purpose of this paper is to determine the a -invariant, the Hilbert function of the ring A/I_X , a set of generators for the ideal I_X , as well as the minimum distance of the code $C_X(d)$ when X is the set of K -rational points of the Segre variety defined over the finite field K .

2. GENERAL RESULTS

In this section, we recall some basic definitions such as the Hilbert function, the a -invariant of an ideal, the Segre variety and general results that will be useful later on.

Let K , $X \subseteq \mathbf{P}_l(K)$ with $|X| = t$, A and $I = I_X$ be as defined in the previous section and let $R = A/I$ be the coordinate ring of X . The *Hilbert function* of R is defined as $H_X(d) = \dim_K A_d - \dim_K I_d$, (cf. [1, Chapter 9, Section 3]).

Let $I = \bigoplus_{i=\gamma}^{\infty} I_i$, where γ is the lowest degree of a nontrivial homogeneous component of the ideal I . The a -invariant of R (or the a -invariant of I or even the a -invariant of X) is the integer a_X such that

- (a) $H_X(d) = \dim_K A_d = \binom{l+d}{d} \Leftrightarrow d < \gamma$;
- (b) $H_X(d) < H_X(d+1) < t$ if $0 \leq d < a_X$;
- (c) $H_X(d) = t$ if $d > a_X$.

The number $a_X + 1$ is called the *regularity index* of R by some authors.

For any field L let $\mathbf{P}_r(L)$ be the r -dimensional projective space over the field L . Recall that the Segre map is defined as [6 p. 25, 7, Section 25.5]

$$\begin{aligned}\varphi: \mathbf{P}_n(L) \times \mathbf{P}_m(L) &\rightarrow \mathbf{P}_N(L), \\ \varphi(\underline{a}, \underline{b}) &= (a_0b_0, \dots, a_ib_j, \dots, a_nb_m),\end{aligned}$$

where $\underline{a} = (a_0, \dots, a_n)$, $\underline{b} = (b_0, \dots, b_m)$ and $N = (n+1)(m+1) - 1$.

The image of this map is known as the Segre variety defined over the field L and it will be denoted by S . If $Z_{ij} = X_iY_j$ this variety is the locus of the quadratic polynomials $Z_{ij}Z_{kl} - Z_{il}Z_{kj}$.

From now on, an element of $\mathbf{P}_N(L)$ will be denoted by $\underline{z} = (z_{00}, \dots, z_{nm})$ and let $A_n = L[X_0, \dots, X_n]$, $A_m = L[Y_0, \dots, Y_m]$ and $A_N = L[Z_{00}, \dots, Z_{nm}]$.

In the sequel, L will denote the finite field K with $q = p^r$ elements (p prime and r a positive integer) and $S = S_K$ the set of K -rational points of the Segre variety defined over K .

Note that, since the Segre map φ is an embedding,

$$\#(S_K) = \#(\mathbf{P}_n(K))\#(\mathbf{P}_m(K)) = \left(\frac{q^{n+1}-1}{q-1}\right)\left(\frac{q^{m+1}-1}{q-1}\right).$$

Let \bar{K} be the algebraic closure of K and \bar{X} be the image of the map $\bar{\varphi}: \mathbf{P}_n(\bar{K}) \times \mathbf{P}_m(\bar{K}) \rightarrow \mathbf{P}_N(\bar{K})$, given by $\bar{\varphi}(\underline{a}, \underline{b}) = (a_0b_0, \dots, a_ib_j, \dots, a_nb_m)$. If $A_N = K[Z_{00}, \dots, Z_{nm}]$ let $\bar{A}_N = \bar{K}[Z_{00}, \dots, Z_{nm}]$ and let $I_{\bar{X}}$ be the vanishing ideal of \bar{X} in \bar{A}_N .

Remark 2.1. Since \bar{K} is algebraically closed $I_{\bar{X}} = \langle I_{\bar{X}}(2) \rangle$, (cf. [6, p. 51]).

Remark 2.2. The following result, which will be useful later, was proved in [9] (see also [10]): $I_{\mathbf{P}_n(K)} = \langle X_i^q X_j - X_i X_j^q : 0 \leq i < j \leq n \rangle$.

3. THE HILBERT FUNCTION OF S_K

Let K be a finite field with q elements and let $S = S_K$ be the set of K -rational points of the Segre variety as introduced above. For any $I = (i_0, \dots, i_n) \in \mathbf{N}^{n+1}$, X^I will denote the monomial $X_0^{i_0} \cdots X_n^{i_n}$. Let $B = K[X_0Y_0, \dots, X_nY_m] = \bigoplus_{d \geq 0} B_d$ be the graded subalgebra of $K[X_0, \dots, X_n, Y_0, \dots, Y_m]$ with the grading given by

$$B_d = \left\{ \sum_{I,J} a_{I,J} X^I Y^J : a_{I,J} \in K, |I| = |J| = d \right\}, \quad d \geq 0.$$

LEMMA 3.1. *For $d \geq 2$ the kernel of the surjective linear transformation $\theta: A_N(d) \rightarrow B_d$, $f \rightarrow \theta_f$, where $\theta_f(\underline{X}, \underline{Y}) = f(X_0 Y_0, \dots, X_n Y_n)$, is $I_S(2)A_N(d-2)$.*

Proof. It is easy to see that $I_S(2)A_N(d-2) \subset \ker \theta$. Obviously, $\ker \theta \subset I_{\bar{S}} = \langle Q_{ij} \rangle$ with $Q_{ij} \in A_N$ being quadrics. By using the generalized division algorithm, any $f \in \ker \theta$ can be written as $f = \sum_{i,j} \lambda_{ij} Q_{ij} + r$, where $\lambda_{ij} \in A_N(d-2)$ and $r \in A_N$ is not divisible by $LT(Q_{ij})$, the leading term of Q_{ij} . Since $\{Q_{ij}\}$ forms a Gröbner basis for $I_{\bar{S}}$ it follows that $r = 0$ (since $f \in I_{\bar{S}}$) and the claim is proved. ■

Let $V_d = I_{\mathbf{P}_n}(d)A_m(d) + A_n(d)I_{\mathbf{P}_m}(d) \subset B_d$. The following result will be useful in determining the Hilbert function of the ring A_N/I_S .

PROPOSITION 3.1. *For $d \geq 2$ the kernel of the surjective linear transformation $\pi \circ \theta$ given by*

$$A_N(d) \xrightarrow{\theta} B_d \xrightarrow{\pi} B_d/V_d$$

$$f \rightarrow \theta_f \rightarrow \theta_f + V_d$$

is $I_S(d)$.

Proof. If $f \in \ker(\pi \circ \theta)$ then $\theta_f \in V_d$. In this case $\theta_f = \sum_i (h_i m_i(Y) + M_i(X) h'_i)$ where $h_i \in I_{\mathbf{P}_n}(d)$, $m_i(Y) \in A_m(d)$, $M_i(X) \in A_n(d)$ and $h'_i \in I_{\mathbf{P}_m}(d)$. Therefore $f(\varphi(P, Q)) = \theta_f(P, Q) = 0 \ \forall P \in \mathbf{P}_n(K)$, $Q \in \mathbf{P}_m(K)$, and it follows that $f \in I_S(d)$.

If $f \in I_S(d)$ then $\theta_f(P, Q) = 0 \ \forall P \in \mathbf{P}_n(K)$, $Q \in \mathbf{P}_m(K)$, i.e., $\theta_f \in I_{\mathbf{P}_s(K)}$ with $s = m + n + 1$. The following cases arise:

(1) $d < q + 1$, $2d < q + 1$. Using Remark 2.2 above and the fact that $\theta_f \in B_d$, it follows that $\theta_f \equiv 0$ and therefore $f \in V_d$.

(2) $d < q + 1$, $2d \geq q + 1$. Since $\theta_f \in B_d$ and B_d consists of bihomogeneous polynomials of bidegree (d, d) , it follows that $\theta_f \equiv 0$ and therefore $f \in V_d$.

(3) $d \geq q + 1$. In this case,

$$\theta_f = \sum_u m_u(\underline{X}, \underline{Y})(X_i^q X_j - X_i X_j^q) + \sum_v M_v(\underline{X}, \underline{Y})(Y_i^q Y_j - Y_i Y_j^q)$$

where $m_u(\underline{X}, \underline{Y})$ and $M_v(\underline{X}, \underline{Y})$ are monomials of degree $2d - (q + 1)$. However,

$$\theta_f = \sum_u m'_u(\underline{X})(X_i^q X_j - X_i X_j^q) m''_u(\underline{Y}) + \sum_v M'_v(\underline{Y})(Y_i^q Y_j - Y_i Y_j^q) M''_v(\underline{X})$$

with $m''_u(\underline{Y}) \in A_m(d)$ and $M''_v(\underline{X}) \in A_n(d)$. Thus $\theta_f \in V_d$. ■

COROLLARY 3.1. *The Hilbert function of the ring A_N/I_S is given by*

$$H_S(d) = H_{\mathbf{P}_n(K)}(d)H_{\mathbf{P}_m(K)}(d).$$

Proof. From Proposition 3.1 it follows that $A_N(d)/I_S(d) \simeq B_d/V_d$. Moreover, $B_d/V_d \simeq A_n(d)/I_{\mathbf{P}_n(K)}(d) \otimes_K A_m(d)/I_{\mathbf{P}_m(K)}(d)$, (the natural isomorphism). By taking dimensions the claim follows. ■

Remark 3.1. From the above corollary it follows that

$$\dim_K C_S(d) = H_{\mathbf{P}_n(K)}(d)H_{\mathbf{P}_m(K)}(d).$$

COROLLARY 3.2. *The a -invariant of the ring A_N/I_S is given by*

$$a_S = \max\{n(q-1), m(q-1)\}.$$

Proof. It is an immediate consequence of the previous corollary and the fact that $a_{\mathbf{P}_n(K)} = n(q-1)$ and $a_{\mathbf{P}_m(K)} = m(q-1)$ (cf. [10]). ■

4. THE VANISHING IDEAL OF S_K

The following result provides a set of generators for the vanishing ideal of the set S of K -rational points of the Segre variety defined over the finite field K with q elements.

THEOREM 4.1. *The vanishing ideal of the set S of K -rational points of the Segre variety is given by*

$$I_S = \langle I_S(2), I_S(q+1) \rangle.$$

Proof. In the proof of Proposition 3.1 it was noted that if $f \in I_S(d)$ for $d < q+1$, then $\theta_f \equiv 0$. Therefore, $f \in I_S(2)A_N(d-2)$, i.e., $I_S(d) \subset \langle I_S(2) \rangle$ whenever $d < q+1$.

If $d > q+1$, in order to prove the theorem, it is enough to show that $I_S(d) \subset \sum_{i,j} Z_{ij}I_S(d-1)$. In this case, with the notation as in Proposition 3.1, we observe that if $m_u(\underline{X}, \underline{Y}) = X_0^{i_0} \cdots X_n^{i_n} Y_0^{j_0} \cdots Y_m^{j_m}$ and $M_v(\underline{X}, \underline{Y}) = X_0^{a_0} \cdots X_n^{a_n} Y_0^{b_0} \cdots Y_m^{b_m}$, then $\sum_{r=0}^n i_r > 0$, $\sum_{r=0}^m j_r > 0$, $\sum_{r=0}^n a_r > 0$ and $\sum_{r=0}^m b_r > 0$. Therefore, $\theta_f = \sum_{i,j} X_i Y_j h_{ij}(\underline{X}, \underline{Y})$ with $h_{ij}(\underline{X}, \underline{Y}) \in I_{\mathbf{P}_s(K)}(2d-2)$. Thus $f = \sum_{i,j} Z_{ij} \theta_{ij}(\underline{Z})$ with $\theta_{ij}(\underline{Z}) \in I_S(d-1)$, showing that $f \in \sum_{i,j} Z_{ij} I_S(d-1)$. ■

5. THE MINIMUM DISTANCE OF THE CODE $C_S(d)$

Let K be a finite field with q elements and let $S = S_K$ be the set of K -rational points of the Segre variety. In this section the minimum distance of the Reed–Muller-type code $C_S(d)$ is determined.

Recall that the minimum distance of the projective Reed–Muller code $C_{\mathbf{P}_l(K)}(d)$ is $\delta_{\mathbf{P}_l}(d) = (q - s)q^{l-r-1}$, where $d - 1 = r(q - 1) + s$, $0 \leq s < q - 1$ (cf. [14]).

THEOREM 5.1. *With the same notation as above, the minimum distance $\delta_S(d)$ of the code $C_S(d)$ is given by*

$$\delta_S(d) = \delta_{\mathbf{P}_n}(d)\delta_{\mathbf{P}_m}(d).$$

Proof. Let $f \in K[Z_{00}, \dots, Z_{nm}]_d$ and let $S = \{P_{11}, \dots, P_{k_1 k_2}\}$ where $k_1 = |\mathbf{P}_n(K)|$ and $k_2 = |\mathbf{P}_m(K)|$. Let $\Lambda = (f(P_{11}), \dots, f(P_{k_1 k_2})) \in C_S(d) - \{0\}$. Note that if φ is the Segre map, then for all i, j , $P_i \in \mathbf{P}_n(K)$ and $Q_j \in \mathbf{P}_m(K)$ exist such that $\varphi(P_i, Q_j) = P_{ij}$, i.e., $(X_0 Y_0, \dots, X_n Y_m)(P_i, Q_j) = P_{rs}$. Therefore,

$$\Lambda = (f(X_0 Y_0, \dots, X_n Y_m)(P_1, Q_1), \dots, f(X_0 Y_0, \dots, X_n Y_m)(P_{k_1}, Q_{k_2})).$$

Let $f(X_0 Y_0, \dots, X_n Y_m) = \sum_{i,j} a_{I,J} X^I Y^J$. For each $P \in \mathbf{P}_n(K)$ and $Q \in \mathbf{P}_m(K)$ let $f_P(Y) = \sum a_{I,J} P^I Y^J \in A_m(d)$ and $f_Q(X) = \sum a_{I,J} X^I Q^J \in A_n(d)$, respectively.

Let $\Lambda_i = (f_{P_i}(Q_1), \dots, f_{P_i}(Q_{k_2})) \in C_{\mathbf{P}_m}(d)$, $i = 1, \dots, k_1$, and let $k_3 = \#\{i : \Lambda_i \neq 0\} > 0$. Since the Hamming weight of Λ_i is such that $wt(\Lambda_i) \geq \delta_{\mathbf{P}_m}(d)$ for all i with $\Lambda_i \neq 0$, it follows that $wt(\Lambda) \geq k_3 \delta_{\mathbf{P}_m}(d)$.

In a similar way, for each $j \in \{1, \dots, k_2\}$ let $\Gamma_j = (f_{Q_j}(P_1), \dots, f_{Q_j}(P_{k_1})) \in C_{\mathbf{P}_n}(d)$. Take j such that $\Gamma_j \neq 0$. If $k_3 < \delta_{\mathbf{P}_n}(d)$ then $wt(\Gamma_j) \leq k_3 < \delta_{\mathbf{P}_n}(d)$. Thus $wt(\Lambda) \geq \delta_{\mathbf{P}_n}(d)\delta_{\mathbf{P}_m}(d)$.

Let $\Omega_1 = (g(P_1), \dots, g(P_{k_1})) \in C_{\mathbf{P}_n}(d)$ with $wt(\Omega_1) = \delta_{\mathbf{P}_n}(d)$ and $g \in A_n(d)$, and similarly let $\Omega_2 = (h(Q_1), \dots, h(Q_{k_2})) \in C_{\mathbf{P}_m}(d)$ with $wt(\Omega_2) = \delta_{\mathbf{P}_m}(d)$ and $h \in A_m(d)$. Since the linear transformation θ (as defined in Proposition 3.1) is surjective, $F \in A_N(d)$ exists such that

$$(gh(P_1, Q_1), \dots, gh(P_{k_1}, Q_{k_2})) = (F(\varphi(P_1, Q_1)), \dots, F(\varphi(P_{k_1}, Q_{k_2}))) =$$

$\Omega \in C_S(d)$ and $wt(\Omega) = \delta_{\mathbf{P}_n}(d)\delta_{\mathbf{P}_m}(d)$, proving the Theorem. ■

6. AN EXAMPLE

In order to illustrate the main results given above, an example is provided.

Let $K = GF(4) = \{0, 1, a, a^2\}$ be the field with four elements and let $\varphi: \mathbf{P}_1(K) \times \mathbf{P}_1(K) \rightarrow \mathbf{P}_3(K)$ be the Segre embedding over K . It is easy to see that the set of K -rational points of the Segre variety is the following:

$$\begin{aligned} S = \{ & (0, 0, 0, 1), (0, 0, 1, 0), (0, 1, 0, 0), (1, 0, 0, 0), (0, 0, 1, a), (0, 0, 1, a^2), \\ & (0, 0, 1, 1), (0, 1, 0, a), (1, a, a, a^2), (1, 0, a, 0), (1, a^2, a, 1), (1, 1, a, a), \\ & (1, a, 0, 0), (1, a^2, 0, 0), (1, 1, 0, 0), (0, 1, 0, a^2), (1, a, a^2, 1), (1, 0, a^2, 0), \\ & (1, a^2, a^2, a), (1, 1, a^2, a^2), (0, 1, 0, 1), (1, a, 1, a), (1, 0, 1, 0), (1, a^2, 1, a^2), \\ & (1, 1, 1, 1)\}. \end{aligned}$$

Since $a_{\mathbf{P}_1(K)} = 3$, it follows from Corollary 3.2 that $a_S = 3$. Now take $d = 2$. Then from Corollary 3.1, $\dim_K C_S(2) = H_S(2) = 9$, from Theorem 5.1 the minimal distance of the code is 9, and from Theorem 4.1, $I_S = \langle I_S(2), I_S(5) \rangle$. If $Z_{00}, Z_{10}, Z_{01}, Z_{11}$ denote the coordinates of $\mathbf{P}_3(K)$, a set of generators for I_S is

$$\begin{aligned} & \{Z_{10}Z_{01} - Z_{00}Z_{11}, Z_{01}^4Z_{11} - Z_{01}Z_{11}^4, Z_{00}Z_{01}^3Z_{11} - Z_{00}Z_{11}^4, Z_{00}^2Z_{01}^2Z_{11} \\ & - Z_{00}Z_{10}Z_{11}^3, Z_{00}^3Z_{01}Z_{11} - Z_{00}Z_{10}^2Z_{11}^2, Z_{10}^4Z_{11} - Z_{10}Z_{11}^4, Z_{00}Z_{10}^3Z_{11} \\ & - Z_{00}Z_{11}^4, Z_{00}^2Z_{10}^2Z_{11} - Z_{00}Z_{01}Z_{11}^3, Z_{00}^3Z_{10}Z_{11} - Z_{00}Z_{01}^2Z_{11}^2, Z_{00}^4Z_{11} \\ & - Z_{00}Z_{11}^4, Z_{00}^4Z_{01} - Z_{00}Z_{11}^4, Z_{00}^4Z_{10} - Z_{00}Z_{11}^4\}. \end{aligned}$$

Now, if $Z_{ij} = X_iY_j$ for all $i = 0, 1, j = 0, 1$, we observe that the generators for I_S can be obtained from the elements of the K -linear space $V = [I_{\mathbf{P}_1(K)}(2) \otimes_K A_1(2)] + [A_1(2) \otimes_K I_{\mathbf{P}_1(K)}(2)]$.

For instance, if we take $H = Y_0Y_1^4(X_0^4X_1 - X_0X_1^4) \in V$, we note that:

$$H = (X_0Y_0)(X_0^3Y_1^3)(X_1Y_1) - X_0Y_0(X_1Y_1)^4 = Z_{00}Z_{01}^3Z_{11} - Z_{00}Z_{11}^4 \in I_S.$$

Also, a basis $\{f_1, f_2, \dots, f_9\}$ for the vector space $A_3(2)/I_S(2)$ can be given: $f_1 = Z_{00}^2, f_2 = Z_{00}Z_{01}, f_3 = Z_{00}Z_{10}, f_4 = Z_{00}Z_{11}, f_5 = Z_{01}^2, f_6 = Z_{01}Z_{11}, f_7 = Z_{10}^2, f_8 = Z_{10}Z_{11}, f_9 = Z_{11}^2$, from which a generating matrix for the code $C_S(2)$ can be obtained. The Macaulay System [3] was used in the calculations of this example.

ACKNOWLEDGMENT

The authors would like to thank the referee for the helpful suggestions which greatly improved the presentation of the paper.

REFERENCES

1. D. Cox, J. Little, and D. O'Shea, "Ideals, Varieties and Algorithms," UTM, Springer-Verlag, Berlin, 1992.
2. I. Duursma, C. Rentería, and H. Tapia-Recillas, Reed-Muller codes on complete intersections, in *Applicable Algebra in Engineering, Communication and Computing*, AAEECC, Vol. 11, pp. 455–462, Springer-Verlag, Berlin (2001).
3. D. R. Grayson and M. Stillman, *Macaulay*, Vol. 2, 1998, <http://www.math.uiuc.edu/Macaulay2/>.
4. J. P. Hansen, Zero-dimensional schemes, in "Proceedings of International Conference, Ravello, 1992" (F. Orrechia and L. Chiantini, Eds.), Walter de Gruyter, Berlin, 1994.
5. J. P. Hansen, Linkage and codes on complete intersections, preprint.
6. J. Harris, "Algebraic Geometry: A First Course," GTM No. 133, Springer-Verlag, Berlin, 1992.
7. J. W. P. Hirschfeld and J. A. Thas, "General Galois Geometries," Oxford Univ. Press, Oxford, 1991.
8. R. F. Lax and G. Salazar, An analogue of one-point codes for complete intersection varieties, preprint.
9. D. J. Mercier and R. Rolland, Polynômes homogènes à plusieurs variables sur un corps fini \mathbb{F}_q qui s'annulent sur l'espace projectif $\mathbb{P}_m(\mathbb{F}_q)$, *J. Pure Appl. Algebra* **124** (1998), 227–240.
10. C. Rentería and H. Tapia-Recillas, Reed-Muller codes: An ideal theory approach. *Comm. Algebra* **25** (1997), 401–413.
11. C. Rentería and H. Tapia-Recillas, The a -invariant of some Reed-Muller codes, in *Applicable Algebra in Engineering, Communications and Computing (AAEECC)*, Vol. 10, No. 1, pp. 33–40, Springer-Verlag, Berlin (1999).
12. C. Rentería and H. Tapia-Recillas, Reed-Muller type codes on the Veronese variety over finite fields, in "Coding Theory, Cryptography and Related Areas" (J. Buchmann, T. Hoholdt, H. Stichtenoth, and H. Tapia-Recillas, Eds.), pp. 237–243, Springer-Verlag, Berlin, 2000, ISBN 3-540-66248-0.
13. H. G. Schaathun, The weight hierarchy of product codes, *IEEE Trans. Inform. Theory* **46** (2000), 2648–2651.
14. A. B. Sørensen. Projective Reed-Muller codes. *IEEE Trans. on Inform. Theory* **37** (1991), 1567–1576.
15. V. K. Wei and K. Yang, On the generalized Hamming weights of product codes. *IEEE Trans. Inform. Theory* **39** (1993), 1709–1713.